



## IMPROVING THE QUALITY OF THE LIFE STATUS OF INFANTS WITH UNILATERAL CONGENITAL CLEFT LIP AND PALATE

**Mirzaeva Firuza Avazovna**

Bukhara State Medical Institute Assistant of the Department of Surgical Dentistry  
[feruza\\_zar@mail.ru](mailto:feruza_zar@mail.ru)

**Yadgarova Gulnora Sadritdinovna**

PhD of the Department of Surgical Dentistry, Bukhara State Medical Institute.  
[farhodik2010@mail.ru](mailto:farhodik2010@mail.ru)

**Annotation:** The most urgent medical and social problem is congenital malformations, due to their high frequency and severity. This is a significant problem for public health and justifies the need to continue studying the prevalence of this pathology in various regions. Every year, for every 100 thousand of the population, the number of newborns with RGN increases by 1.38 (Isakov.L.O., 2009). According to WHO, the frequency of birth of children with CCLP in the world is 0.6-1.6 cases per 1000 newborns. Congenital cleft lip and palate is one of the most common malformations of the maxillofacial area with a tendency to increase (Hong M. et al., 2017; Inoyatov.A.Sh., 2020). Congenital cleft lip and palate in combination with other defects is one of the causes of perinatal mortality and comes out on top in the structure of childhood morbidity and disability, representing not only a medical but also an important social problem. According to many authors, the number of children with this pathology is constantly increasing and in the next decade the frequency of such cases will be 2 times higher than 100 years ago.

**Key words:** cleft, children, lip, palate, silicone, obturator.

## УЛУЧШЕНИЕ КАЧЕСТВА ЖИЗНИ НОВОРОЖДЕННЫХ С ОДНОСТОРОННЕЙ ВРОЖДЕННОЙ РАСЩЕЛИНОЙ ГУБЫ И НЕБА

**Мирзаева Феруза Авазовна**

ассистент, Бухарский государственный медицинский институт  
[feruza\\_zar@mail.ru](mailto:feruza_zar@mail.ru)

**Ядгарова Гулнора Садритдиновна**

PhD, Бухарский государственный медицинский институт  
[farhodik2010@mail.ru](mailto:farhodik2010@mail.ru)

**Аннотация:** Наиболее актуальной медико-социальной проблемой являются врожденные пороки развития, обусловленные их высокой частотой и тяжестью. Это является серьезной проблемой для общественного здравоохранения и обосновывает необходимость



продолжения изучения распространенности этой патологии в различных регионах. Ежегодно на каждые 100 тысяч населения число новорожденных с РГН увеличивается на 1,38 (Исаков Л.О., 2009). По данным ВОЗ, частота рождения детей с ССЛР в мире составляет 0,6-1,6 случая на 1000 новорожденных. Врожденная расщелина губы и неба является одним из наиболее распространенных пороков развития челюстно-лицевой области с тенденцией к увеличению (Hong M. et al., 2017; Inoyatov.A.Sh., 2020). Врожденная расщелина губы и неба в сочетании с другими дефектами является одной из причин перинатальной смертности и занимает первое место в структуре детской заболеваемости и инвалидности, представляя собой не только медицинскую, но и важную социальную проблему. По мнению многих авторов, число детей с этой патологией постоянно растет и в ближайшее десятилетие частота таких случаев будет в 2 раза выше, чем 100 лет назад.

**Ключевые слова:** расщелина, дети, губа, нёба, силикон, obturator.

Cleft lip occurs twice as often in boys as in girls. A cleft palate is twice as common in girls.

The frequency of births of children with CCLP ranges from 1 in 500 to 1 in 1000, depending on the place and time of the research (Ushnitsky I.D., et al., 2015; Mamedov A.A. et al., 2017; Arsenina. O.I., 2017). It should be noted that in the regions of the Russian Federation the frequency of developmental pathologies varies widely: from 1:653 to 1:1280, which may be due to the characteristics of the regions and the characteristics of populations.

Barillas (2009). B. Bergendal et al. (2008.2010), C.I.Emeka et al. (2017) showed a trend towards a steady increase in the incidence of VA, and in recent years, the proportion of cleft palate among the defects has increased. The system of complex treatment of children with VA provides for a multi-stage interdisciplinary interaction of specialists.

One out of five newborns with a cleft of the MFR has a syndromic pathology (Blokhina S.I., 1992). Thus, in the Republic of Belarus, every fifth inhabitant suffered from the Chernobyl accident, and there is an annual increase in the number of children with congenital maxillofacial anomalies by 0.25 times per 1000 newborns. A.S. Artyushkevich et al. (2004) indicate that in the Republic of Belarus over the past 10 years, according to genetic monitoring, there has been an increase in the frequency of birth of children with this pathology.

In the article by A.Sh. Inoyatov et al. (2016) conducted an epidemiological study to study the spread of CVLP in the Navoi region (Uzbekistan, Bukhara) for the period from 2005 to 2009. 136 births of children with CCLP were identified, the average incidence of malformation was 1 chance every 620 births. A continuous growth in dynamics (by years) of the number of CVLPs was noted. In the Navoi region and the city of Navoi, there is a high frequency of CVLP. According to the USBC (United States Bureau of Census), the birth rate of children with congenital AFA in the world is on average 1:600. According to the statistical data of Kazakhstan, the frequency of birth of children with congenital malformations of the MFR in the republic is high and amounts to 1:880. More than 6,000 children are registered with the dispensary, about 400 children are born annually.

In the Republic of Bashkortostan, the birth rate of children with CCLP is higher than the average for the Russian Federation (Topolnitsky O.Z., Chuikin O.S., 2015). In the Perm region



and the city of Perm, the frequency of congenital pathology of the maxillofacial area among newborns is 1:700 (Danilova M.A., 2006, 2013).

Children with congenital cleft lip and palate need long-term complex and systematic treatment and are registered at the dispensary (Simanovskaya E.Yu., Shcheglova A.P., 2002; Isakov L.O. 2013., 2015).

Congenital cleft lip and palate occupies a leading place in the structure of all antenatal anomalies and leads to a change in the aesthetics of the child's face, a violation in the immune system, nutritional function, respiration, speech defects (Bimbas E.S., et al., 2018; Tokarev P.V., 2015, 2019; Yakubova. Z.Kh., et al., 2020; Willadsen E. 2012; Plsek P., 2013). And as a result, these are children with disabilities from birth until the complete elimination of functional disorders (Egorova M.V. et al., 2010; Azimov M.I., 2012; Persin L.S., 2020).

According to statistics from a number of European countries, the frequency of births of children with cleft lip and palate is one in 500-1000 newborns (Materna Kiryluk A. et al., 2014; Meazzini M.C. et al., 2018). The overall frequency of morphological malformations in children under 1 year of age is approximately 27.2 per 1000 population.

Conclusion: Congenital cleft lip and palate (CCLP) is a severe malformation of the maxillofacial area, accompanied by gross anatomical and functional disorders. Difficulties in restoring impaired vital functions of nutrition, respiration and speech, anatomical restoration of the upper lip and, nose and upper jaw in a growing body are the cause of disability in children with cleft lip and palate for many years.

Treatment of cleft lip and palate is a complex multifaceted and unresolved problem. For optimal treatment results from birth to 14-16 years of age. Obviously, active monitoring of a child with CCLP is necessary from the moment of birth. Given the pronounced anatomical, functional, cosmetic problems, as well as the presence of comorbidities, dynamic monitoring and treatment by various specialists is necessary - an oral surgeon, an otorhinolaryngologist, a dentist, a pediatrician, and a speech therapist. Only the combined efforts of doctors of different profiles can achieve a good result in the treatment of such children.

#### LIST OF LITERATURE

1. A.S. Artyushkevich et al. "Age-related morphology of the sternum". 2013 Minsk.
2. M.A. Pogrel, K.E. Kahnberg, L. Andersson "Essentials of Oral and Maxillofacial Surgery" 2014 Cothenburg.
3. M.E. Zorich, O.S. Yatskevich, A.I. Karanevich, 2013; ON THE. Peleshenko, "The choice of methods of surgical treatment of patients with congenital cleft palate" 2013. Tajikistan.
4. Козин И.А. Эстетическая хирургия врожденных расщелин лица. – М., 1996.
5. Rahmatilloevna K. F. et al. Prevention of Major Dental Diseases in Preschool Children //Eurasian Medical Research Periodical. – 2022. – Т. 6. – С. 172-174.
6. Rahmatilloevna K. F. et al. Improving the Methods of Treatment and Prevention of Complications of Odontogenic Diseases of the Jaws in Children //Eurasian Medical Research Periodical. – 2022. – Т. 6. – С. 168-171.
7. Kamalova F. R., Safarova M. S. ARTIFICIAL FEEDING AND DENTAL HEALTH //湖南大学学报 (自然科学版). – 2021. – Т. 48. – №. 7.
8. Rahmatillaevna K. F., Umarovich S. S. Consequences Of Early Extraction of Deciduous Teeth in Children //Eurasian Medical Research Periodical. – 2021. – Т. 1. – №. 1. – С. 51-54.
9. Камалова Ф. Р. ПЕРВИЧНАЯ ПРОФИЛАКТИКА СТОМАТОЛОГИЧЕСКИХ ЗАБОЛЕВАНИЙ У ДЕТЕЙ СТРАДАЮЩИХ С САХАРНЫМ ДИАБЕТОМ //Новый день в медицине. – 2020. – №. 2. – С. 383-386.



- 
10. RajabovA. A. INDICATIONS FOR THE DISEASE OF INFLAMMATORY DISEASES OF THE DENTAL SYSTEM IN CHILDREN WITH EXAMINED CEREBRAL PALSY //EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE. – 2022. – T. 2. – №. 3. – С. 56-60.
  11. KAMBAROVA S. A. Effect of Surgical Manipulation in Morphometric Growth of Maxillofacial Area at Children with Congenital Lip and Palate Splits At I and Ii Period of Childhood // Annals of the Romanian Society for Cell Biology. – 2021. - Vol. 25. - Issue 4. – P. 1853 – 1858.
  12. Kambarova Sh.A. IDENTIFICATION OF THE MORPHOMETRIC PARAMETERS OF THE CRANIO-FASCIAL REGION OF CHILDREN WITH CONGENITAL CLEFT AND PALATE REFLECTIONS USING A DEVELOPED RESEARCH MAP // CENTRAL ASIAN JOURNAL OF MEDICAL AND. – 2021. - Vol. 2. - Issue 3. – P. 286 – 290.
  13. Камбарова С.А. ВЛИЯНИЕ ХИРУРГИЧЕСКОЙ МАНИПУЛЯЦИИ НА МОРФОМЕТРИЧЕСКОЕ РАЗВИТИЕ ЛИЦА И ЧЕЛЮСТИ У ПАЦИЕНТОВ С ВРОЖДЕННЫМИ РАССТОЯНИЯМИ ГУБ И ЧАСТНОГО ДВИГАТЕЛЯ // Новый день в медицине. – 2021. - P. 128 - 130.
  14. Kambarova Sh.A., Pulatova Sh.K. Revitalization of nonspecific immunity factors in patients with diffuse phlegmoine of the maxillo facial area using Bakteriofags // New day in medicine. - 2020. - P. 128 - 130.